

Mathematics

ON ONE SPECTRUM OF UNIVERSALITY FOR WALSH SYSTEM

M. A. NALBANDYAN*

Chair of Higher Mathematics (Department of Physics) YSU, Armenia

In the present work it is shown that the set $D = \left\{ \sum_{i=0}^{\infty} \delta_i 2^{N_i} : \delta_i = 0, 1 \right\}$ for every sequence $N_0 < N_1 < \dots < N_i < \dots$ of natural numbers can be changed into the set of the form $\Lambda = \{k + o(\omega(k)) : k \in D\}$, where $\omega(k)$ is an arbitrary, tending to infinity at $k \rightarrow +\infty$ sequence, such that Λ is the spectrum of universality for Walsh system.

Keywords: Walsh system, universal series, representation theorems, representations by subsystems.

Introduction. Let S be a space of functions defined on $[0, 1]$ (for example, $S = L^p[0, 1]$) and let T be a type of convergence (for example, the convergence in $L^p[0, 1]$ metric or the almost everywhere convergence). Here we will mainly consider $S = L^0[0, 1]$ – the class of all almost everywhere finite, measurable functions and $T =$ almost everywhere convergence on $[0, 1]$.

A series

$$\sum_{k=1}^{\infty} a_k \varphi_k(x) \tag{1}$$

is said to be *universal in the usual sense* for S, T , if for any function $f(x) \in S$ there exists an increasing sequence of natural numbers n_k , such that the corresponding sequence of partial sums $\sum_{j=1}^{n_k} a_j \varphi_j(x)$ converges to $f(x)$ in the sense of T .

There are also other types of universality such as *universality with respect to rearrangements* for S, T : the latter means that for any function $f(x) \in S$ there exists rearrangement $k \mapsto \sigma(k)$ such that the series $\sum_{k=1}^{\infty} a_{\sigma(k)} \varphi_{\sigma(k)}(x)$ converges to $f(x)$ in the sense of T .

We will also say that the series (1) is universal in the sense of partial series

* E-mail: nalbandyanmikayel@yahoo.com

for S, T , if for any function $f(x) \in S$ there exists a partial series $\sum_{k=1}^{\infty} a_{n_k} \varphi_{n_k}(x)$ of (1), which converges to $f(x)$ in the sense of T .

The first example of trigonometric series universal in the usual sense for the class of all measurable functions has been constructed by D.E. Menshoeff [1] (see also [2]). This result was extended by A.A. Talalian [3] to arbitrary complete orthonormal systems. He also established [4], that if $\{\varphi_n(x)\}_{n=1}^{\infty}, x \in [0,1]$, is an arbitrary orthonormal system, then there exist a series $\sum a_k \varphi_k(x)$, which is universal in the sense of partial series for the class of all measurable functions and T =convergence in measure on $[0,1]$. The following general result was obtained by M. Grigorian [5]:

Theorem. The class of orthogonal series simultaneously possessing the following properties 1), 2) are not empty:

1) universality with respect to rearrangements and in the sense of partial series both in each $L^p[0,1], p \in [1,2)$, and in $\bigcap_{1 \leq p < 2} L^p[0,1]$;

2) universality with respect to rearrangements and in the sense of partial series for S =all measurable functions and T =almost everywhere convergence on $[0,1]$.

The fact that there exists a functional series universal with respect to rearrangements for S =class of almost everywhere finite, measurable functions and T =almost everywhere convergence, was mentioned by W. Orlicz [6]. Note that Riemann has proved (see [7], p. 317) that every unconditionally convergent numerical series is universal with respect to rearrangements for S = all reals.

Definition. The set of natural numbers Λ , for which it is possible to construct an universal (in some sense) series $\sum_{\lambda_k \in \Lambda} a_k \varphi_{\lambda_k}(x)$, we will call the spectrum of universality (in the same sense).

In the rest of the paper we will consider universal series in Walsh system.

Let $\omega(k)$ be an arbitrary sequence, tending to infinity as $k \rightarrow +\infty$. By the small change of some set D we will mean the set $\{k + o(\omega(k)) : k \in D\}$.

Such small transformations of sets were considered for the first time by G. Kozma and A. Olevskii [8], with the aim to transform these sets into representation spectrum. More precisely, it was proved by them for trigonometric system that for any sequence $w(k)$ tending to infinity there is a symmetric representation spectrum $\Lambda = \{\pm k^2 + o(w(k))\}_{k \in \mathbb{N}}$, i.e. each measurable function f allows the representation $f(x) = \sum_{n \in \Lambda} c_n(f) e^{inx}$, where the sum converges almost everywhere.

This result was extended to the Walsh system by the author in [9], namely:

Theorem. For arbitrary $l \in \{2^k\}_{k=0}^{\infty}$ there exists a subsystem $\{w_{n_k}\}_{k=1}^{\infty}, n_k \in \{k^l + o(k^{l-1})\}_{k \in \mathbb{N}}$ of Walsh system such that for every measurable function there exists a series by subsystem $\{w_{n_k}\}_{k=1}^{\infty}$ converging a.e. to this function. In other words, there exists a representation spectrum of the form $\Lambda_l = \{k^l + o(k^{l-1})\}_{k \in \mathbb{N}}, l \in \{2^k\}_{k=0}^{\infty}$.

Theorem. For arbitrary sequence $\{\omega(k)\}_{k=1}^{\infty}$, tending to infinity, there exists a subsystem $\{w_{n_k}\}_{k=1}^{\infty}$, $n_k \in \{k^2 + o(\omega(k))\}_{k \in N}$, of Walsh system such that for arbitrary measurable function there exists a series by subsystem $\{w_{n_k}\}_{k=1}^{\infty}$ converging a.e. to this function, i.e. there exists a representation spectrum $\Lambda = \{k^2 + o(\omega(k))\}_{k \in N}$ (the notation $\{k^2 + o(\omega(k))\}_{k \in N}$ means that we can find a sequence $\alpha_k \rightarrow 0$ such that $\{k^2 + \alpha_k \cdot \omega(k)\}_{k \in N}$ is a representations spectrum).

Let us consider the set of natural numbers in binary representation: $N = \left\{ \sum_{i=0}^{\infty} \delta_i 2^i : \delta_i = 0, 1 \right\}$. After substituting all indexes i in the exponents by N_i (for a given sequence $N_0 < N_1 < \dots < N_i < \dots$) we will get the set $D = \left\{ \sum_{i=0}^{\infty} \delta_i 2^{N_i} : \delta_i = 0, 1 \right\}$, which, as it can be easily seen, cannot be a universality spectrum in general. However, for any sequence $\omega(k)$, tending to infinity, by small change of D it can be transformed into a spectrum of universality for the Walsh system. The main result of the present work is the following

Theorem. For any sequence of nonnegative integers $N_0 < N_1 < \dots < N_i < \dots$ and arbitrary sequence $\omega(k)$, tending to infinity, the set $D = \left\{ \sum_{i=0}^{\infty} \delta_i 2^{N_i} : \delta_i = 0, 1 \right\}$ can be transformed into the set $\Lambda = \{k + o(\omega(k)) : k \in D\} = \{\lambda_n\}_{n=1}^{\infty}$ by small change such that Λ is a universality spectrum (in $S = L^0[0, 1]$ and in the sense of T -convergence almost everywhere) for Walsh system, i.e. there exists a series $\sum_{k=1}^{\infty} a_k w_{\lambda_k}(x)$ with $a_i \rightarrow 0$, such that for arbitrary function $f \in L^0[0, 1]$ there is a sequence of natural numbers $\{\nu_k\}$ such that $\lim_{k \rightarrow \infty} \sum_{i=1}^{\nu_k} a_i w_{\lambda_i}(x) = f(x)$ almost everywhere on $[0, 1]$.

Definitions, Notations and Some Properties. Let us recall the definition of Walsh system $\{w_k(t)\}_{k=0}^{\infty}$ in the Paley ordering [10, 11]:

$$w_0(t) = 1, \quad w_1(t) = \begin{cases} 1, & t \in [0, 1/2), \\ -1, & t \in [1/2, 1], \end{cases} \quad w_{2^k}(t) = w_1(2^k t),$$

and for natural q with binary representation $q = \sum_{i=0}^{\infty} q_i 2^i$, where $q_i = 0$ or $q_i = 1$,

we define $w_q(t) = \prod_{i=0}^{\infty} (w_{2^i}(t))^{q_i}$. Using this definition, it is easy to check the following properties, which we will use later in the text:

- 1) for every natural number q we have $w_q(2^k t) = w_{q \cdot 2^k}(t)$;
- 2) if natural numbers p and q have nonintersecting binary representation

(see definition below), then $w_p(t)w_q(t) = w_{p+q}(t)$ (the property of index addition).

Let $p = 2^{i_0} + \dots + 2^{i_k}$ and $q = 2^{j_0} + \dots + 2^{j_n}$ be some natural numbers. We will say that binary representations of numbers p and q do not intersect, if $\{i_0, \dots, i_k\} \cap \{j_0, \dots, j_n\} = \emptyset$.

Let $f(t) \in L[0,1]$ and $\hat{f}(k) = \int_0^1 f(t)w_k(t)dt$ be its Fourier–Walsh coefficient. Then for each polynomial $P(t)$ in Walsh system we have:

$$P(t) = \sum_{k \geq 0} \hat{P}(k)w_k(t); \quad (*)$$

1. $(P)_m = \sum_{k=0}^m \hat{P}(k)w_k$;
2. $\text{spec}\{P\}$ represents the set of those nonnegative integers k , for which w_k appears in the representation (*);
3. $\text{deg}\{P\}$ is the maximal element of $\text{spec}\{P\}$;
4. $\|\hat{P}\|_1 = \sum_{k \in \text{spec}\{P\}} |\hat{P}(k)|$.

The Construction of the Spectrum of Universality. For the given sequence $N = \{N_0, N_1, \dots, N_k, \dots\}$ of increasing nonnegative integers we define the following sets:

$$S(i, n) = \left\{ \sum_{k=0}^n \delta_k 2^{N_k^{(i,n)}} : \delta_k = 0, 1; N_k^{(i,n)} \in N \right\} \text{ and } B_n = \bigcup_{i=0}^n (i + S(i, n)),$$

where $N_k^{(i,n)}$ are chosen such that the following conditions are satisfied:

1. $\frac{n}{\omega(\min\{S(i, n)\})} < \frac{1}{n}$ for all $0 \leq i \leq n$;
2. $\max\{S(i-1, n)\} < \min\{S(i, n)\}$, $1 \leq i \leq n$;
3. $\max\{B_{n-1}\} < \min\{B_n\}$.

Then, for sufficiently large n , we have $B_n = \{k + o(\omega(k)) : k \in D\}$. Note that $S = \bigcup_{n=0}^{\infty} \bigcup_{i=0}^n (S(i, n)) \subset D$ and small change of it a subset D is specified. Other elements of D will be changed by 0, which is also a special case of small change.

Thus, $\Lambda' = \bigcup_{n=0}^{\infty} B_n = \{k_m + o(\omega(k_m)) : k_m \in D\} = \{\lambda_n\}_{n=1}^{\infty} \subset \{k + o(\omega(k)) : k \in D\} = \Lambda$.

We will prove that Λ' is a universality spectrum, which means that Λ is a universality spectrum too. To prove that Λ' is a universality spectrum, it is enough to prove the following lemma.

Main Lemma. For every $f \in L^0[0,1]$ and for arbitrary $\varepsilon > 0$, $\delta > 0$ and $k_0 \in N$ there exists a polynomial $P(x)$ in Walsh system such that:

1. $P(x) = \sum_{k=k_0}^{\bar{k}} a_k w_{\lambda_k}(x)$;
2. $\lambda_k \in \Lambda$;

3. $|a_k| < \delta$;
4. $\text{mes}\{|f(x) - P(x)| > \delta\} < \varepsilon$.

Proof of the Main Lemma. First we need to prove the following lemma.

Lemma. For any $|a| < 1$, $0 < \alpha < 1$, $y > 0$ and any $N_i \in \mathbb{N}$ with

$N_0 < N_1 < \dots < N_{k-1}$ there exists a polynomial $W(t) = \sum_{i=1}^{2^k-1} \hat{W}(i) \bar{w}_i(t)$ such that:

1. $m\{t : |1 - W(t)| \geq y\} < y^{-\alpha} c^k$;
2. $|\hat{W}(i)| \leq a$,

where $c = \frac{(1-a)^\alpha + (1+a)^\alpha}{2} < 1$ and $\bar{w}_i(t) = w_{q_0 2^{N_0} + \dots + q_{k-1} 2^{N_{k-1}}}(t)$ for $i = q_0 2^0 + \dots + q_{k-1} 2^{k-1}$, $q_j = 0, 1$, $0 \leq j \leq k-1$.

In the rest of the paper to emphasize that the polynomial $W(t)$ in the Lemma depends on numbers N_0, N_1, \dots, N_{k-1} , we will denote $W(t) = W(t) \{N_0, \dots, N_{k-1}\}$.

Proof. For the natural numbers $N_0 < N_1 < \dots < N_{k-1}$ we denote $\varphi_m(t) = a \cdot w_1(2^{N_{m-1}} t) = a \cdot w_{2^{N_{m-1}}}(t)$ with $|a| < 1$, then $\varphi_k = a$ on the first half of each interval $\Delta_i^{(k)} = \left[\frac{i-1}{2^{N_{k-1}}}, \frac{i}{2^{N_{k-1}}} \right]$, $1 \leq i \leq 2^{N_{k-1}}$, and $\varphi_k = -a$ on the second half.

Now for $\alpha < 1$ we have

$$\int_{\Delta_i^{(k)}} (1 - \varphi_k(t))^\alpha dt = \frac{|\Delta_i^{(k)}|}{2} ((1-a)^\alpha + (1+a)^\alpha) = c \int_{\Delta_i^{(k)}} dt,$$

where we denote $c = \frac{(1-a)^\alpha + (1+a)^\alpha}{2} < \left(\frac{1-a+1+a}{2} \right)^\alpha = 1$.

It is easy to see that φ_j , for $0 \leq j < k-1$, are constant on each of $\Delta_i^{(k)}$, $1 \leq i \leq 2^{N_{k-1}}$. Let us prove that $\int_0^1 (1 - \varphi_1(t))^\alpha \dots (1 - \varphi_n(t))^\alpha dt = c^n$.

For $n=1$ it is obvious. Let us assume that the statement is true for $n = k-2$ and prove it for $n = k-1$. We have

$$\begin{aligned} \int_0^1 (1 - \varphi_1(t))^\alpha \dots (1 - \varphi_{k-1}(t))^\alpha dt &= \sum_{i=1}^{2^{N_{k-1}}} \int_{\Delta_i^{(k)}} (1 - \varphi_1(t))^\alpha \dots (1 - \varphi_{k-2}(t))^\alpha (1 - \varphi_{k-1}(t))^\alpha dt = \\ &= \sum_{i=1}^{2^{N_{k-1}}} (1 - \varphi_1(t_i))^\alpha \dots (1 - \varphi_{k-2}(t_i))^\alpha \int_{\Delta_i^{(k)}} (1 - \varphi_{k-1}(t))^\alpha dt, \end{aligned}$$

where $t_i \in \Delta_i^{(k)}$. Then

$$\begin{aligned} \int_0^1 (1 - \varphi_1(t))^\alpha \dots (1 - \varphi_{k-1}(t))^\alpha dt &= c \cdot \sum_{i=1}^{2^{N_{k-1}}} (1 - \varphi_1(t_i))^\alpha \dots (1 - \varphi_{k-2}(t_i))^\alpha \int_{\Delta_i^{(k)}} dt = \\ &= c \cdot \int_0^1 (1 - \varphi_1(t))^\alpha \dots (1 - \varphi_{k-1}(t))^\alpha dt = c \cdot c^{k-1} = c^k \end{aligned}$$

Now we present the product $(1 - \varphi_0(t)) \cdots (1 - \varphi_k(t))$ in the form of the sum:

$$(1 - a w_1(2^{N_0} t)) \cdots (1 - a w_1(2^{N_{k-1}} t)) = \sum_{i=0}^{2^k-1} \hat{w}(i) \bar{w}_i(t),$$

where for each $i = q_1 2^0 + \cdots + q_k 2^{k-1}$, $q_j = 0, 1$ we denote $\bar{w}_i(t) = w_{q_0 2^{N_0} + \cdots + q_{k-1} 2^{N_{k-1}}}(t)$. It is easy to see that $\hat{w}(0) = 1$ and $|\hat{w}(i)| \leq a$ for $0 < i < 2^k$. Thus, for nonintersecting m and n we have $\bar{w}_m \cdot \bar{w}_n = \bar{w}_{m+n}$. By denoting $W(t) = -\sum_{i=1}^{2^k-1} \hat{w}(i) \bar{w}_i(t)$, we have $\int_0^1 |1 - W(t)|^\alpha dt < c^k$ and, therefore, $m\{t : |1 - W(t)| \geq y\} \leq y^{-\alpha} \int_0^1 |1 - W(t)|^\alpha dt < y^{-\alpha} \cdot c^k$.

The second statement of the Lemma is obvious from the construction of the polynomial.

The Lemma is proved.

Proof of the Main Lemma. Let us approximate the function f by polynomial P_1 so that $m\{t : |P_1 - f| > \delta/2\} < \varepsilon/2$. We take a such that $0 < a < \delta$,

n such that $(\deg P_1 + 1) \frac{2^\alpha \|\hat{P}_1\|_1^\alpha}{\delta^\alpha} c^n < \frac{\varepsilon}{2}$ and take $y = \frac{\delta}{2 \|\hat{P}_1\|_1}$.

We define the polynomial $P(t) = \sum_{k=0}^{\deg P_1} \hat{P}_1(k) w_k(t) W_k(t)$, where the polynomials $W_k(t) = W(t) \{N_1^{(k)}, \dots, N_n^{(k)}\}$ are chosen according to the Lemma, and the numbers $N_k^{(i)}$ are to be chosen later.

Now we put $M = \max\{\deg P_1, n\}$. For all $m \geq M$ we can choose the numbers $N_k^{(i)}$ from the set of numbers $N_k^{(i,m)}$ such that $\text{spec}\{P\} \subset B_m$ for all $m \geq M$ and, therefore, $\text{spec}\{P\} \subset \{\lambda_k\}_{k=1}^\infty$. Hence, we can choose numbers $N_k^{(i)}$ such that $\min\{\text{spec}\{P\}\} > k_0$ for any given k_0 . So the first and second statements of the Main Lemma are satisfied.

We have the following estimates: $|P - P_1| = \left| \sum_{k=0}^{\deg P_1} \hat{P}_1(k) w_k(t) (W_k(t) - 1) \right|,$

$$m\left\{t : |P - P_1| \geq \sum_{k=0}^{\deg P_1} |\hat{P}_1(k)| y\right\} \leq m\left\{t : \sum_{k=0}^{\deg P_1} |\hat{P}_1(k) (W_k(t) - 1)| \geq \sum_{k=0}^{\deg P_1} |\hat{P}_1(k)| y\right\} \leq \sum_{k=0}^{\deg P_1} m\left\{t : |\hat{P}_1(k) (W_k(t) - 1)| \geq |\hat{P}_1(k)| y\right\} \leq \sum_{k=0}^{\deg P_1} m\left\{t : |W_k(t) - 1| \geq y\right\} \leq (\deg P_1 + 1) y^{-\alpha} c^n.$$

Then $m\{t : |P - f| > \delta/2 + \|\hat{P}_1\|_1 y\} \leq m\{t : |P - P_1| + |P_1 - f| > \delta/2 + \|\hat{P}_1\|_1 y\} \leq m\{t : |P - P_1| > \|\hat{P}_1\|_1 y\} + m\{t : |P_1 - f| > \delta/2\} < (\deg P_1 + 1) y^{-\alpha} c^n + \varepsilon/2 < \varepsilon$.

So, we have $m\{t : |P - f| > \delta\} < \varepsilon$.

The Main Lemma is proved.

Proof of the Theorem.

Theorem. There exists a series $\sum_{k=1}^{\infty} c_k w_{\lambda_k}(x)$ with $c_k \rightarrow 0$, which is universal in the usual sense for $L^0[0,1]$.

Proof. We denote by $\{f_n(x)\}_{n=1}^{\infty}$ the sequence of polynomials with rational coefficients and, applying successively the Main Lemma, we can choose a sequence of polynomials $Q_j(x)$ in subsystem of the Walsh system

$Q_j(x) = \sum_{i=m_{j-1}}^{m_j-1} a_i w_{\lambda_i}(x)$, satisfying the following conditions:

1. $m \{x : |f_k(x) - \sum_{j=1}^k Q_j(x)| < 2^{-k}\} > 1 - 2^{-k}$;
2. $|a_i| < 2^{-j}$, for all $i \in [m_{j-1}, m_j)$.

Let $f(x) \in L^0[0,1]$. Let us choose a subsequence of polynomials $\{f_{v_k}\}$ such that $m \{x : |f(x) - f_{v_k}(x)| < 2^{-2k}\} > 1 - 2^{-k}$. Let $B_k = \{x : |f(x) - f_{v_k}(x)| < 2^{-2k}\}$,

$E_k = \left\{x : \left|f_{v_k} - \sum_{j=1}^{v_k} Q_j(x)\right| < 2^{-v_k}\right\}$ and, finally, $E = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} (E_k \cap B_k)$. Obviously,

$|E| = 1$. Then $\left|f(x) - \sum_{j=1}^{v_k} \left(\sum_{i=m_{j-1}}^{m_j-1} a_i w_{\lambda_i}(x)\right)\right| < 2^{-k}$ for all $x \in E_k \cap B_k$.

This means that $\lim_{k \rightarrow \infty} \sum_{i=1}^{v_k} a_i w_{\lambda_i}(x) = f(x)$ on E , i.e. $\sum_{i=1}^{\infty} a_i w_{\lambda_i}(x)$ is universal in the usual sense for $L^0[0,1]$ and $a_i \rightarrow 0$.

The author is grateful to prof. M. Grigoryan for useful remarks and discussions.

Received 10.05.2011

REFERENCES

1. **Menshoff D.E.** // Mat. Sbornik, 1947, v. 20, p. 197 (in Russian).
2. **Kozlov W.Ya.** // Mat. Sbornik, 1950, v. 26, p. 351 (in Russian).
3. **Talalyan A.A.** // Izv. AN Arm. SSR. Matematika, 1957, v. 10, № 3, p. 17 (in Russian).
4. **Talalyan A.A.** // Uspehi Mat. Nauk, 1960, v. 15, № 5, p. 567–604 (in Russian).
5. **Grigorian M.G.** // Izv. NAN Armenii. Matematika, 2000, v.35, № 4, p. 23–29 (in Russian).
6. **Orlicz W.** // Bull. de l'Academie Polonaise des Sciences, 1927, v. 81, p. 117–125.
7. **Fichtengolz G.M.** A Course of Differential and Integral Calculus. V.II. M.: Nauka, 1996 (in Russian).
8. **Kozma G., Olevskii A.** // J. Anal. Math., 2001, v. 84, p. 361–393.
9. **Nalbandyan M.A.** // Izv. Vuzov. Matematika, 2009, v. 10, p. 51–62 (in Russian).
10. **Walsh J.L.** // Amer. J. Math., 1923, v. 45, p. 5–24.
11. **Paley R.E.A.C.** // Proc. London Math. Soc., 1932, v. 34, p. 241–279.

Մ.Ա. Նալբանդյան. Ուոլշի համակարգի ունիվերսալության, որոշակի սպեկտրի մասին

Աշխատանքում ցույց է տրված, որ ցանկացած բնական թվերի համար բազմությունը փոքր փոփոխությամբ կարելի է դարձնել բազմություն, որտեղ – կամայական անվերջի ձգտող հաջորդականություն է, երբ , որ լինի ունիվերսալության սպեկտր:

М.А. Налбандян. Об одном спектре универсальности для системы Уолша

В работе показано, что для любых натуральных $N_0 < N_1 < \dots < N_i < \dots$ множество $D = \left\{ \sum_{i=0}^{\infty} \delta_i 2^{N_i} : \delta_i = 0, 1 \right\}$ малым изменением можно так превратить в множество $\Lambda = \left\{ k + o(\omega(k)) : k \in D \right\}$, где $\omega(k)$ – произвольная, стремящаяся к бесконечности последовательность при $k \rightarrow +\infty$, что Λ будет спектром универсальности.